

«
3, 49008, ; -mail: edgladky@gmail.com»

()

(),

)

When solving parametric reliability problems, one often has to construct distributions of statistical data to find the probability of containment in the operability region. This paper considers the problem of 2D statistical ensemble fitting. The use of a 2D normal distribution in statistical data description is not always justified because statistical ensembles rather frequently (at the level of marginal components and a stochastic relationship between them) have properties different from the normal case. From a practical standpoint, it is desirable for researchers to describe 2D statistical ensembles with the use of universal distributions, which allow one to cover a wide range of source data using a single analytical form. In the process of fitting, account should be made of bounded ranges of random variables.

The paper considers who universal distribution construction methods, which are based on 1D orthogonal Jacobi polynomial expansions. In these distributions, the random variable range is a rectangle. In the first method, a 2D distribution is constructed using a direct expansion in the 1D Jacobi polynomials. A 2D Jacobi distribution function and regression lines are obtained, and methods to fit it are considered. In theory, a distribution obtained in this way can be used, up to the fourth order inclusive, for marginal and even reduced moments different from the normal case. However, its real capabilities are limited to values of reduced moments (1D and even) that differ from the normal case only very slightly. Otherwise, the probability surface may enter negative ranges with the occurrence of multiple modes.

The second way to construct a 2D distribution is to use a normal copula and 1D Jacobi distributions as components. The resulting 2D distribution allows one to deal with 1D distributions different from the normal case and linear correlation. This approach is justified because, according to research data, it is a linear stochastic relationship that relates a significant part of 2D statistical ensembles, and marginal distributions deviate from the normal case. Regression lines of a distribution of this kind are obtained, and it is shown that they are curved because marginal distributions differ from the normal one. The paper considers the practical example of fitting a 2D

© . . . , . . . , 2022

. - 2022. - 4.

ensemble of characteristics of a liquid-propellant rocket engine some components of which are related via a linear stochastic relationship (the parameters that characterize a nonlinear stochastic relationship proved to be insignificant) and have 1D distributions different from the normal one. The fitted and observed frequencies are in rather good agreement. It is shown that a distribution based on a normal copula is more universal, and it is recommended for practical calculations.

Keywords: 2D distribution, orthogonal Jacobi polynomials, Jacobi distribution, normal copula, fitting.

() , () .

$$(, , , [1])$$

(), () .

$$(-) .$$

[7].

[10].

$$f(x) = \sum_{k=0}^4 c_k P_k(x; r, s, a, b) h(x; r, s, a, b), \quad (1)$$

$h(x; r, s, a, b)$ –
 $P_k(x; r, s, a, b)$ –

$k; c_k$ –

$[a, b]$

$$h(x; r, s, a, b) = (x-a)^r (b-x)^s \quad (r > -1 \quad s > -1).$$

$$P_k(x; r, s, a, b) \quad [2],$$

$$P_k(x; r, s, a, b) = \sum_{i=0}^k \tilde{b}_k^{(i)} x^i, \quad (2)$$

$\tilde{b}_k^{(i)}$ –

$$\tilde{b}_k^{(i)} = \sum_{j=i}^k b_k^{(j)} (b-a)^{k-j} 2^{i-k} C_j^i (-b-a)^{j-i}.$$

$b_k^{(i)}$

i

$P_k(x; r, s, -1, 1)$,

$[-1, 1]$.

[2, 8].

c_k (1)

$$c_k = \frac{1}{\|P_k(x; r, s, a, b)\|^2} \sum_{i=0}^k \tilde{b}_k^{(i)} r_i,$$

r_i –

(

).

$$ja(x) = c_0 h(x; r, s, a, b) \sum_{i=0}^4 b_{sum}^{(i)} x^i, \quad (3)$$


$$c_0 = \frac{\Gamma(r+s+2)}{(b-a)^{r+s+1} \Gamma(r+1) \Gamma(s+1)}, \quad b_{sum}^{(i)} = \frac{1}{c_0} \sum_{j=0}^{4-i} \tilde{b}_{4-j}^{(i)} c_{4-j}.$$

(1)

$$JA(x) = I_{\bar{x}}(\Gamma + 1, S + 1) + \sum_{i=1}^4 c_i P_{i-1}(x; \Gamma + 1, S + 1, a, b) h(x; \Gamma + 1, S + 1, a, b), \quad (4)$$

$$I_t(p, q) = \frac{\Gamma(p+q)}{\Gamma(p)\Gamma(q)} \int_0^t x^{p-1} (1-x)^{q-1} dx - \quad ; \quad \bar{x} = \frac{x-a}{b-a}.$$

» () « ,
 ($m = 0, \dagger = 1$) [2] . 1. -
 . 1 -
 () .



. 1 -

$$ja_1(\bar{x}) = \prod_{i=1}^2 h_i(x_i; \bar{n}_i) \sum_{0 \leq k_1 + k_2 \leq 4} c_{k_1 k_2} P_{k_1}(x_1; \bar{n}_1) P_{k_2}(x_2; \bar{n}_2), \quad (5)$$

$$\Omega = \{a_1 \leq x_1 \leq b_1; a_2 \leq x_2 \leq b_2\}$$

$$P_{k_i}(x_i; \bar{n}_i) - \quad k_i -$$

$$h_i(x_i; \bar{n}_i); \quad \bar{n}_i = \{\Gamma_i, S_i, a_i, b_i\} -$$

; $c_{k_1 k_2} -$

$$c_{k_1 k_2} = \frac{1}{\prod_{i=1}^2 \|P_{k_i}(x_i; \bar{n}_i)\|^2} \iint_{\Omega} P_{k_1}(x_1; \bar{n}_1) P_{k_2}(x_2; \bar{n}_2) f(\bar{x}) dx_1 dx_2. \quad (6)$$

(5)

((2))

$$P_{k_1}(x_1; \bar{n}_1) = \sum_{l_1=0}^{k_1} b_{k_1}^{(l_1)} x_1^{l_1}; \quad P_{k_2}(x_2; \bar{n}_2) = \sum_{l_2=0}^{k_2} b_{k_2}^{(l_2)} x_2^{l_2} .$$

(5)

$$P_{k_1}(x_1; \bar{n}_1) P_{k_2}(x_2; \bar{n}_2) = \sum_{l_1=0}^{k_1} \sum_{l_2=0}^{k_2} b_{k_1}^{(l_1)} b_{k_2}^{(l_2)} x_1^{l_1} x_2^{l_2}$$

(5).

$$\begin{bmatrix} c_{00} P_0(x_1) P_0(x_2) \\ c_{10} P_1(x_1) P_0(x_2) & c_{01} P_0(x_1) P_1(x_2) \\ c_{20} P_2(x_1) P_0(x_2) & c_{11} P_1(x_1) P_1(x_2) & c_{02} P_0(x_1) P_2(x_2) \\ c_{30} P_3(x_1) P_0(x_2) & c_{21} P_2(x_1) P_1(x_2) & c_{12} P_1(x_1) P_2(x_2) & c_{03} P_0(x_1) P_3(x_2) \\ c_{40} P_4(x_1) P_0(x_2) & c_{31} P_3(x_1) P_1(x_2) & c_{22} P_2(x_1) P_2(x_2) & c_{13} P_1(x_1) P_3(x_2) & c_{04} P_0(x_1) P_4(x_2) \end{bmatrix}$$

(5)

$$\begin{aligned} & x_1^0 x_2^0, \\ & ; \\ & c_{11} P_1(x_1) P_1(x_2), \end{aligned}$$

(5)

$$x_1^1 x_2^1,$$

$$\sum_{0 \leq k_1 + k_2 \leq 4} c_{k_1 k_2} P_{k_1}(x_1; \bar{n}_1) P_{k_2}(x_2; \bar{n}_2) = \sum_{0 \leq k_1 + k_2 \leq 4} b_{k_1 k_2} x_1^{k_1} x_2^{k_2},$$

$$b_{k_1 k_2} = \sum_{\substack{l_1=k_1 \\ l_1+l_2 \leq 4}}^4 \sum_{l_2=k_2}^4 c_{l_1 l_2} b_{l_1}^{(k_1)} b_{l_2}^{(k_2)} .$$

$$j a_1(\bar{x}) = \prod_{i=1}^2 h_i(x_i; \bar{n}_i) \sum_{0 \leq k_1 + k_2 \leq 4} b_{k_1 k_2} x_1^{k_1} x_2^{k_2} . \quad (7)$$

$$(7) \quad k_1 k_2, \quad (6),$$

$$k_1 k_2 = \frac{b_1 b_2 \int \int P_{k_1}(x_1; \bar{n}_1) P_{k_2}(x_2; \bar{n}_2) f(\bar{x}) dx_1 dx_2}{a_1 a_2 \prod_{i=1}^2 \|P_{k_i}(x_i; \bar{n}_i)\|^2} =$$

$$= \frac{\int_{a_1}^{b_1} \int_{a_2}^{b_2} \sum_{l_1=0}^{k_1} \sum_{l_2=0}^{k_2} b_{k_1}^{(l_1)} b_{k_2}^{(l_2)} x_1^{l_1} x_2^{l_2} f(\bar{x}) dx_1 dx_2}{\prod_{i=1}^2 \|P_{k_i}(x_i; \bar{n}_i)\|^2} = \frac{\sum_{l_1=0}^{k_1} \sum_{l_2=0}^{k_2} b_{k_1}^{(l_1)} b_{k_2}^{(l_2)} r_{l_1, l_2}}{\prod_{i=1}^2 \|P_{k_i}(x_i; \bar{n}_i)\|^2}, \quad (8)$$

$$0 \leq k_1 + k_2 \leq 4; \quad r_{l_1, l_2} - \quad l_1, l_2.$$

$$, \quad m_2(x_1)$$

$$m_2(x_1) = \frac{\int_{a_2}^{b_2} x_2 \prod_{i=1}^2 h(x_i; \bar{n}_i) \sum_{0 \leq k_1 + k_2 \leq 4} \sum c_{k_1 k_2} P_{k_1}(x_1; \bar{n}_1) P_{k_2}(x_2; \bar{n}_2) dx_2}{\sum_{k_1=0}^4 c_{k_1} P_{k_1}(x_1; \bar{n}_1) h(x_1; \bar{n}_1)} =$$

$$= \frac{\sum_{0 \leq k_1 + k_2 \leq 4} \sum c_{k_1 k_2} P_{k_1}(x_1; \bar{n}_1) \int_{a_2}^{b_2} x_2 P_{k_2}(x_2; \bar{n}_2) h(x_2; \bar{n}_2) dx_2}{\sum_{k_1=0}^4 c_{k_1} P_{k_1}(x_1; \bar{n}_1)}.$$

$$, \quad k_2 \neq 1.$$

$$m_2(x_1) = \frac{\|P_1(x_2; \bar{n}_2)\|^2 \sum_{k_1=0}^3 c_{k_1, 1} P_{k_1}(x_1; \bar{n}_1)}{b_2^{(1)} \cdot \sum_{k_1=0}^4 c_{k_1} P_{k_1}(x_1; \bar{n}_1)} \quad (a_1 \leq x_1 \leq b_1).$$

$$m_2(x_1)$$

$$m_1(x_2)$$

$$(7),$$

$$(5),$$

$$JA_1(t_1, t_2) = \int_{a_1}^{t_1} \int_{a_2}^{t_2} ja_1(x_1, x_2) dx_1 dx_2 = I_{\bar{t}_1}^-(r_1 + 1, s_1 + 1) I_{\bar{t}_2}^-(r_2 + 1, s_2 + 1) +$$

$$+ I_{\bar{t}_1}^-(r_1 + 1, s_1 + 1) \sum_{k_2=1}^4 c_{k_2} P_{k_2-1}(t_2; r_2 + 1, s_2 + 1, a_1, b_1) h(t_2; r_2 + 1, s_2 + 1, a_2, b_2) +$$

$$+ I_{\bar{t}_2}^-(r_2 + 1, s_2 + 1) \sum_{k_1=1}^4 c_{k_1} P_{k_1-1}(t_1; r_1 + 1, s_1 + 1, a_1, b_1) h(t_1; r_1 + 1, s_1 + 1, a_1, b_1) +$$

$$+ \sum_{\substack{2 \leq k_1 + k_2 \leq 4 \\ k_1, k_2 \neq 0}} c_{k_1 k_2} P_{k_1-1}(t_1; r_1 + 1, s_1 + 1, a_1, b_1) h(t_1; r_1 + 1, s_1 + 1, a_1, b_1) \times$$

$$\times P_{k_2-1}(t_2; \Gamma_2 + 1, S_2 + 1, a_2, b_2) h(t_2; \Gamma_2 + 1, S_2 + 1, a_2, b_2),$$

$$\bar{t}_i = \frac{t_i - a_i}{b_i - a_i} \quad (i = 1, 2).$$



$$JA_1(t_1, t_2),$$

$$: m_1 = m_2 = 0, \tau_1 = \tau_2 = 1, \dots_{11} = 0.2, \beta_{1_1} = \beta_{1_2} = 0.1,$$

$$\beta_{2_1} = \beta_{2_2} = 0.2, \dots_{12} = \dots_{21} = 0.1, \dots_{13} = \dots_{31} = 0.7, \dots_{22} = 1.1.$$

$$[-4.0, 4.8]$$

$$\Gamma_1 = S_1 = 9.2, \Gamma_2 = S_2 = 9.2.$$



[2]).

(7).

$$\bar{u}_i = \{\Gamma_i, S_i, a_i, b_i\} \quad (i = 1, 2)$$

$$\Gamma_i = S_i \quad (i = 1, 2).$$

$a_i, b_i.$

(7)

$X_1 O X_2,$

(7)

$$F_{1,2}(x_1, x_2) = C(F_1(x_1), F_2(x_2)),$$

$$C(u, v) \quad (0 \leq u, v \leq 1) - \quad ; F_1(x_1) \quad F_2(x_2) - \\ X_1 \quad X_2.$$

$$\pm 1. \quad [11]$$

$$C_N(u_1, u_2; n) = \frac{1}{2f\sqrt{1-n^2}} \int_{-\infty}^{\Phi^{(-)}(u_1)} \int_{-\infty}^{\Phi^{(-)}(u_2)} \exp\left\{-\frac{y_1^2 - 2ny_1y_2 + y_2^2}{2(1-n^2)}\right\} dy_1 dy_2,$$

$$\begin{aligned} & \quad ; Y_1, Y_2 - \\ & \quad ; n - \quad , \quad Y_1 \quad Y_2; u_1, u_2 - \\ & \quad , \quad X_1 \quad X_2 \quad (u_i = F_i(x_i) \quad i = 1, 2). \end{aligned}$$

$$F_{1,2}(x_1, x_2) = \Phi\left(\Phi^{(-)}(F_1(x_1)), \Phi^{(-)}(F_2(x_2)); n\right), \quad (9)$$

$$\begin{aligned} & (\bullet, \bullet; n) - \quad ; \\ & , \quad (9), \end{aligned}$$

$$f_{1,2}(x_1, x_2) = \frac{N\left(\Phi^{(-)}(F_1(x_1)), \Phi^{(-)}(F_2(x_2)); n\right)}{N\left(\Phi^{(-)}(F_1(x_1))\right) \cdot N\left(\Phi^{(-)}(F_2(x_2))\right)} f_1(x_1) f_2(x_2), \quad (10)$$

$$N(\bullet, \bullet; n), N(\bullet) - \quad ; f_1(x_1), f_2(x_2) - \quad X_1 \quad X_2.$$

$$[9] \quad (\quad [3]).$$

$$(10)$$

$$f_{1,2}(x_1, x_2) = \frac{N\left(\Phi^{(-)}(JA_1(x_1)), \Phi^{(-)}(JA_2(x_2)); n\right)}{N\left(\Phi^{(-)}(JA_1(x_1))\right) \cdot N\left(\Phi^{(-)}(JA_2(x_2))\right)} ja_1(x_1) ja_2(x_2), \quad (11)$$

$$a_i \quad x_i \quad b_i(i=1,2),$$

$$j_{A_1}(x_1), j_{A_2}(x_2) - \dots \quad (3);$$

$$X_1, X_2 \dots$$

$$\Psi(\dots) - m_{X_1} m_{X_2} = \dots \quad (12)$$

$$m_{X_1}, m_{X_2}, \dots \quad (11),$$



$$\Psi(\dots) = M[X_1 \cdot X_2] = \int_{a_1}^{b_1} \int_{a_2}^{b_2} x_1 x_2 f_{1,2}(x_1, x_2) dx_1 dx_2 .$$

$$(12) \dots (11)$$

$$: m_1 = m_2 = 0, \quad \dots \quad (\dots)$$

MathCAD.

$$\dots = 0.72129.$$



$$m_2(x_1) \quad (11)$$

$$m_2(x_1) = \frac{1}{\sqrt{2f} \sqrt{1-\rho^2}} \int_{-\infty}^{\infty} JA_2^{(-)}[\Phi(y_2)] \cdot \exp\left\{-\frac{1}{2}\left(\frac{y_2 - \rho \Phi^{(-)}(JA_1(x_1))}{\sqrt{1-\rho^2}}\right)^2\right\} dy_2 =$$

$$= \int_{-\infty}^{\infty} JA_2^{(-)}[\Phi(y_2)] \cdot N\left(y_2; \rho \Phi^{(-)}(JA_1(x_1)), \sqrt{1-\rho^2}\right) dy_2 ,$$

$$JA_2^{(-)}(\bullet) - X_2 (-) .$$

$$m_1(x_2) = \int_{-\infty}^{\infty} JA_1^{(-)}[\Phi(y_1)] \cdot N\left(y_1; \rho \Phi^{(-)}(JA_2(x_2)), \sqrt{1-\rho^2}\right) dy_1 .$$

(11)

[6].

30 %

$$(12)$$

(X1)

(X2),
n = 148

1.

2 3.

| X_1 | X_2 | X_1 | X_2 | X_1 | X_2 | X_1 | X_2 |
|-------|-------|-------|-------|-------|-------|-------|-------|
| 2.47 | 333 | 2.55 | 329.6 | 2.65 | 328.5 | 2.6 | 326.5 |
| 2.51 | 334 | 2.56 | 331 | 2.57 | 329.2 | 2.55 | 333.1 |
| 2.6 | 332.7 | 2.55 | 329.4 | 2.51 | 332.7 | 2.56 | 330 |
| 2.59 | 333.2 | 2.59 | 330.3 | 2.5 | 333.2 | 2.59 | 331.7 |
| 2.62 | 331.6 | 2.52 | 331.9 | 2.55 | 332.4 | 2.57 | 329 |
| 2.57 | 331 | 2.52 | 331 | 2.5 | 332.8 | 2.52 | 328.3 |
| 2.56 | 331.6 | 2.54 | 330 | 2.55 | 331.7 | 2.56 | 332.5 |
| 2.63 | 325.4 | 2.57 | 326.5 | 2.53 | 333.1 | 2.53 | 332.8 |
| 2.57 | 328.3 | 2.53 | 330.6 | 2.54 | 333.2 | 2.55 | 332.2 |
| 2.49 | 329.6 | 2.51 | 329.8 | 2.62 | 332.3 | 2.53 | 332.5 |
| 2.63 | 331.2 | 2.57 | 330.8 | 2.55 | 331.6 | 2.53 | 331.2 |
| 2.57 | 326.2 | 2.46 | 332.5 | 2.55 | 334.2 | 2.51 | 333.2 |
| 2.57 | 331.9 | 2.56 | 330.8 | 2.62 | 331.6 | 2.52 | 331 |
| 2.58 | 328 | 2.54 | 331.4 | 2.52 | 334.3 | 2.51 | 332.1 |
| 2.48 | 333.8 | 2.51 | 331.3 | 2.49 | 331.5 | 2.52 | 331.6 |
| 2.61 | 327.2 | 2.53 | 332.4 | 2.6 | 331.7 | 2.55 | 331.8 |
| 2.49 | 330.6 | 2.57 | 326.7 | 2.51 | 333 | 2.5 | 330.6 |
| 2.54 | 326.1 | 2.49 | 329.2 | 2.51 | 332.9 | 2.45 | 333.7 |
| 2.49 | 331.4 | 2.58 | 330.3 | 2.6 | 329.1 | 2.58 | 330.7 |
| 2.52 | 331.2 | 2.54 | 332.4 | 2.5 | 331.7 | 2.56 | 332.8 |
| 2.53 | 331.5 | 2.56 | 331.3 | 2.59 | 327.9 | 2.5 | 330.4 |
| 2.54 | 330.5 | 2.55 | 330.8 | 2.5 | 329.8 | 2.56 | 333.4 |
| 2.53 | 329.6 | 2.56 | 328.6 | 2.56 | 330.1 | 2.55 | 330.3 |
| 2.54 | 333.8 | 2.56 | 332.2 | 2.54 | 332.5 | 2.5 | 332.3 |
| 2.53 | 331.5 | 2.64 | 328.8 | 2.56 | 331 | 2.51 | 330.2 |
| 2.55 | 331.1 | 2.6 | 331.3 | 2.55 | 331.9 | 2.55 | 332.4 |
| 2.56 | 331.3 | 2.61 | 330.9 | 2.55 | 332.4 | 2.54 | 332.2 |
| 2.52 | 329.1 | 2.54 | 330.5 | 2.52 | 334 | 2.52 | 330.2 |
| 2.53 | 330.1 | 2.5 | 331.9 | 2.54 | 330 | 2.54 | 332 |
| 2.55 | 327.8 | 2.56 | 330.2 | 2.62 | 328 | 2.5 | 332.7 |
| 2.54 | 330.5 | 2.6 | 330.7 | 2.64 | 326.1 | 2.59 | 328.6 |
| 2.59 | 326.8 | 2.56 | 331.2 | 2.6 | 331.2 | 2.51 | 332.9 |
| 2.53 | 331.7 | 2.61 | 331 | 2.58 | 329.5 | 2.53 | 334.5 |
| 2.53 | 328.5 | 2.54 | 331.7 | 2.57 | 331.1 | 2.56 | 335 |
| 2.56 | 328.7 | 2.49 | 331 | 2.57 | 331 | 2.56 | 330.2 |
| 2.58 | 331.3 | 2.54 | 332.3 | 2.6 | 332.5 | 2.47 | 330.4 |
| 2.5 | 334.7 | 2.57 | 330.2 | 2.6 | 334.2 | 2.54 | 332.8 |

(. 1).
(2 3)
[1].
 $X_2 - [324.0, 336.2]$
« »
($X_1 - [2.43, 2.67]$,
-
(3)

$$ja_1(x_1) = 1,1515 \cdot 10^{-5} \cdot \left(\frac{x_1 - 2,542}{0,025} + 4,478 \right)^{3,16} \left(5,122 - \frac{x_1 - 2,542}{0,025} \right)^{3,16} \times$$

$$\times \left[0,0127 \left(\frac{x_1 - 2,542}{0,025} \right)^4 + 0,0055 \left(\frac{x_1 - 2,542}{0,025} \right)^3 - 0,1188 \left(\frac{x_1 - 2,542}{0,025} \right)^2 - \right.$$

$$\left. - 0,0895 \left(\frac{x_1 - 2,542}{0,025} \right) + 1,1094 \right].$$



$$ja_2(x_2) = 1,8558 \cdot 10^{-6} \cdot \left(\frac{x_2 - 331,763}{1,08} + 7,188 \right)^{3,35} \left(4,108 - \frac{x_2 - 331,763}{1,08} \right)^{3,35} \times$$

$$\times \left[0,0027 \left(\frac{x_2 - 331,763}{1,08} \right)^4 + 0,00115 \left(\frac{x_2 - 331,763}{1,08} \right)^3 - 0,0234 \left(\frac{x_2 - 331,763}{1,08} \right)^2 + \right.$$

$$\left. + 0,3524 \left(\frac{x_2 - 331,763}{1,08} \right) + 1,4795 \right].$$

6.

2 3



. 6 -

1 -

| | X_1 | X_2 |
|----------------|---------|---------|
| (), \bar{X} | 2,55 | 331,0 |
| , s | 0,0392 | 1,94 |
| (), b_1 | 0,1928 | -0,6372 |
| €HT (), b_2 | -0,1320 | 0,3553 |
| , X | 2,542 | 331,763 |

2 -

X_1

| X_1 | 2.453 | 2.478 | 2.504 | 2.529 | 2.555 | 2.581 | 2.606 | 2.632 | 2.657 | Σ |
|-------|------------|------------|--------------|--------------|--------------|--------------|--------------|------------|------------|----------------|
| n_1 | 2 (2.3) | 9 (9.0) | 20 (22.2) | 39 (34.6) | 34 (36.1) | 23 (22.0) | 12 (12.6) | 8 (7.6) | 1 (1.5) | 148 (147.9) |

| | 3- | | | | | | X ₂ | | | | |
|----------------|------------|------------|------------|--------------|--------------|--------------|----------------|--------------|------------|------------|----------------|
| X ₂ | 325.74 | 326.82 | 327.90 | 328.98 | 330.06 | 331.14 | 332.22 | 333.30 | 334.38 | 335.46 | Σ |
| n ₂ | 4 (3.3) | 5 (4.5) | 6 (6.2) | 13 (14.0) | 23 (26.1) | 39 (33.9) | 32 (31.2) | 18 (19.8) | 7 (7.4) | 1 (0.9) | 148 (147.3) |

$$r_{11} = -0.374.$$

$$|r_{11}| > t_{1-\frac{r}{2}} \dagger r_{11}, \quad (13)$$

$$t_{1-\frac{r}{2}} \dagger r_{11} = 1 - \frac{r}{2} (r -$$

$$); \dagger r_{11}^2 = \frac{1}{n} (1 - \dots_{11}^2)^2 + O(1/n^{1.5}).$$

$$r = 0.05 \quad 0.139.$$

r₁₁

$$\begin{aligned} r_{21} &= -0,164; & r_{12} &= 0,340; \\ r_{31} &= -1,142; & r_{13} &= -1,417. \end{aligned}$$

[7],

$$\begin{aligned} \} _{2_1} &= \dots_{21} - S_{1_1 \dots 11}, & \} _{2_2} &= \dots_{12} - S_{1_2 \dots 11}, \\ \} _{3_1} &= \dots_{31} - (S_{2_1} + 3) \dots_{11}, & \} _{3_2} &= \dots_{13} - (S_{2_2} + 3) \dots_{11}. \end{aligned}$$

$$l_{2_1} = -0.092; \quad l_{2_2} = 0.102;$$

$$l_{3_1} = -0.069; \quad l_{3_2} = -0.162.$$

$$l_{2_1}, l_{2_2}, l_{3_1}, l_{3_2}$$

(13)

$$|l_{(\bullet)}| \geq t_{1-\frac{r}{2}} \sqrt{D(l_{(\bullet)})}, \quad (14)$$

$$D(l_2) = \frac{2}{n} (1 - \dots_{11}^2) + O(1/n^{1.5}); \quad D(l_3) = \frac{6}{n} (1 - \dots_{11}^2) + O(1/n^{1.5}).$$

$$(14) \quad l_{2_1} \quad l_{2_2} - 0.211; \quad l_{3_1} \quad l_{3_2} - 0.366$$


$$l_{2_1}, l_{2_2}, l_{3_1}, l_{3_2}$$

X₁ X₂


(12)

MathCAD, " = -0.3163.

(11) (4.)



. 7 - (11)



. 8 - (11)

4

(11),

(11)

| $X_1 \backslash X_2$ | 325.74 | 326.82 | 327.90 | 328.98 | 330.06 | 331.14 | 332.22 | 333.3 | 334.38 | 335.46 | d | |
|----------------------|------------|------------|------------|--------------|--------------|--------------|--------------|--------------|------------|------------|-------|------------------------------|
| 2.453 | | | | (0.1) | (0.2) | – | 1 (0.6) | 1 (0.6) | – | (0.4) | (0.1) | 2 (2.4) |
| 2.478 | | (0.1) | (0.1) | 1 (0.4) | 2 (1.0) | 4 (1.9) | – (2.4) | 2 (2.1) | – | (1.1) | (0.2) | 9 (9.3) |
| 2.504 | (0.2) | (0.3) | (0.5) | – (1.3) | 4 (3.1) | 2 (5.0) | 6 (5.6) | 6 (4.2) | 2 (1.8) | | (0.2) | 20 (22.2) |
| 2.529 | 1 (0.4) | – (0.7) | 1 (1.1) | 2 (2.8) | 8 (5.7) | 9 (8.2) | 10 (8.1) | 5 (5.3) | 3 (1.9) | | (0.2) | 39 (34.4) |
| 2.555 | – (0.8) | – (1.1) | 1 (1.6) | 3 (3.6) | 6 (6.8) | 10 (8.7) | 9 (7.5) | 3 (4.3) | 1 (1.3) | 1 (0.1) | | 34 (35.8) |
| 2.581 | 1 (0.7) | 3 (0.9) | 3 (1.2) | 4 (2.7) | 3 (4.6) | 6 (5.2) | 2 (4) | 1 (2.0) | – (0.6) | | | 23 (21.9) |
| 2.606 | – (0.5) | 2 (0.7) | – (0.9) | 1 (1.7) | – (2.7) | 5 (2.8) | 3 (2.0) | | 1 (0.9) | | | 12 (12.4) |
| 2.632 | 2 (0.5) | – (0.5) | 1 (0.6) | 1 (1.1) | – (1.6) | 3 (1.5) | 1 (1.0) | | (0.4) | (0.1) | | 8 (7.3) |
| 2.657 | (0.1) | (0.1) | (0.2) | 1 (0.3) | (0.3) | (0.3) | (0.1) | | | | | 1 (1.4) |
| d | 4 (3.2) | 5 (4.4) | 6 (6.2) | 13 (14.0) | 23 (26.0) | 39 (34.0) | 32 (31.3) | 18 (19.8) | 7 (7.4) | 1 (0.8) | | 148 (147.1) |

1. , 1975. 399 .
2. , 2017. 1. . 107–122. <https://doi.org/10.15407/itm2017.01.107>
3. , 2018. 4. . 105–118. <https://doi.org/10.15407/itm2018.04.105>
4. , 1966. 588 .
5. , 1975. 648 .
6. , 1971. 576 .
7. , 1997. 2. . 53–65.
8. , 1976. 328 .
9. *Headrick T. C., Mugdadi A.* On simulating multivariate non-normal distributions from the generalized lambda distribution. *Computational statistics & data analysis*. 2006. Volume 50. P. 3343–3353. <https://doi.org/10.1016/j.csda.2005.06.010>
10. *Jonson N. L., Kotz S.* Continuous Multivariate Distributions. Volume 2. N.Y.e.a. John Wiley and Sons, 1972. 333 p.
11. *Lee L.* Generalized econometric models with selectivity. *Econometrica*. 1983. Vol. 51. P. 507–512. <https://doi.org/10.2307/1912003>

01.11.2022,
29.11.2022